



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 173 259 B1**

(12)

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: 06.11.91 (51) Int. Cl.⁵: **A61K 7/06**
(21) Application number: **85110580.9**
(22) Date of filing: **22.08.85**

(54) **Antimicrobial suspensions and antimicrobial hair treatment compositions.**

- (30) Priority: 29.08.84 JP 179978/84
24.12.84 JP 272529/84
27.12.84 JP 276789/84

- (43) Date of publication of application:
05.03.86 Bulletin 86/10

- (45) Publication of the grant of the patent:
06.11.91 Bulletin 91/45

- (94) Designated Contracting States:
AT CH DE FR GB IT LI NL

- (56) References cited:
EP-A- 0 007 704 EP-A- 0 034 385
EP-A- 0 060 611 EP-A- 0 074 819
EP-A- 0 093 541 DE-A- 3 113 872
GB-A- 1 202 716 US-A- 3 489 686
US-A- 3 723 325 US-A- 3 753 916
US-A- 3 940 482

PATENT ABSTRACTS OF JAPAN, unexamined applications, section C, vol. 2, no. 59, April 27, 1978 THE PATENT OFFICE JAPANESE GOVERNMENT page 349 C 78

PATENT ABSTRACTS OF JAPAN, unexamined applications, section C, vol. 2, no. 59, April 27, 1978 THE PATENT OFFICE JAPANESE GOVERNMENT page 349 C 78

NESE GOVERNMENT page 349 C 78

- (73) Proprietor: **Kao Corporation**
14-10, Nihonbashi Kayabacho 1-chome
Chuo-Ku Tokyo 103(JP)

- (72) Inventor: **Takaya, Susumu**
3-20-1, Innai
Funabashi-shi Chiba-ken(JP)
Inventor: **Hirota, Hajime**
5-31-9, Megurohon-cho
Meguro-ku Tokyo(JP)
Inventor: **Nakamura, Motoko**
1-3-34, Motoazabu
Minato-ku Tokyo(JP)

- (74) Representative: **Wächtershäuser, Günter, Dr.**
Tal 29
W-8000 München 2(DE)

EP 0 173 259 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

BACKGROUND OF THE INVENTION

5 I) Field of the Invention

This invention relates to aqueous suspensions of antimicrobial agents and antimicrobial hair treatment compositions in which fine particulate polyvalent metal salts of 2-mercaptopyridine-N-oxide (which may be hereinafter referred to simply as Mept compounds) having a specific size distribution are stably dispersed.

10

II) Description of the Prior Art:

Polyvalent metal salts of 2-mercaptopyridine-N-oxide are known to be effective as antimicrobials and are widely utilized not only as ordinary antimicrobials, but also as anti-dandruff agents and are incorporated into shampoo compositions, hair rinse compositions and the like. The polyvalent metals of the Mept compounds are magnesium, barium, strontium, zinc, cadmium, tin, and zirconium. Of these, zinc salts are widely used.

EP-A-0 093 541 describes anti-dandruff compositions containing zinc pyridine thione (Zpt) in particulate form, wherein a major proportion of the particles has a defined minimum surface area and is in the form of hexagonal flat platelets or fragments thereof. The Zpt particles are incorporated into conventional carriers of a hairdressing, hair conditioner, shampoo, cream, ointment or lotion.

On the other hand, US-patents No. 3,753 916, 3,489,886 and 3,723,325 disclose detergent compositions containing particle deposition and enhancing agents. Certain cationic polymers are used in association with particles having a size ranging from 0.2 to 50 μm to enhance the deposition and retention of such particulate substances on surfaces washed with the detergent composition. The US documents are concerned with the problem of providing detergent compositions, which have improved capacities to impart residual activity or properties to surfaces washed therewith, i.e. to increase the affinity of water-insoluble or sparingly soluble, particular substances for surfaces being washed with the detergent compositions. The US documents do not address the problem of preventing the particulate components of the composition from settling and separating, as time passes.

Mept compounds are sparingly soluble in water, for instance, the zinc salt (hereinafter referred to simply as Zpt compounds) has a solubility in water of 15 ppm at 25° C. Accordingly, when they are incorporated into shampoo or hair rinse compositions, they must be utilized in a form of dispersions.

However, because of the considerable difference between the specific gravities of Zpt compounds (specific gravity = 1.8) and a medium liquid for dispersion, the Zpt compounds are apt to settle and separate as time passes, and thus it was difficult to obtain a stable dispersion system containing Zpt compounds.

In order to prevent such settlement or separation, the following methods are known:

- (i) Make the static viscosity of the dispersion medium high.
- (ii) Make the particle size of Mept compounds so small that Brownian movement will be dominant in the system.

By either method, settlement will be made difficult to take place.

Examples of method (i) are a method of adding viscosity increasing polymers such as cross-linked polyacrylates (Japanese Patent Publication No. 49-49117) and a method of adding acrylic acid/acrylate copolymers (Japanese Patent Publication No. 54-16951). However, these methods have the drawback that limitation is placed on the type of surface active agent usable to stably disperse Mept compounds.

As for method (ii), since Mept compounds having very small particle size are difficult to be produced, compositions containing fine particulate Mept compounds stably dispersed therein have not been practically obtained.

Under such conditions, the present inventors formerly made a study and succeeded in manufacturing Mept compounds having very small particle size (hereinafter referred to as fine particulate Mept compounds) compared with conventional Mept compounds incorporated into shampoo or hair rinse compositions (Japanese Patent Application No. 58-122845, 58-122846 and 59-82690).

The present inventors made further study in order to obtain stable antimicrobial suspensions making use of thus obtained fine particulate Mept compounds, and found that fine particulate Mept compounds are very sensitive to conditions of liquid media for dispersion and are apt to coagulate, leading to difficulty in keeping the original size distribution. More specifically, the following phenomena were noted.

- (1) Fine particulate Mept compounds in a suspension coagulate when electrolytes such as salts are

included in the suspension, or when the suspension is got frozen or heated.

(2) Fine particulate Mept compounds are apt to coagulate when they are incorporated into hair treatment compositions such as shampoos or hair rinses in which surfactants are contained, because the surfactant per se is an electrolyte.

5 In order to prevent the coagulation of fine particulate Mept compounds, surface modification or colloidal protection have been proposed.

Examples of adding a water-soluble polymer to a shampoo composition which incorporates Mept compounds are disclosed in Japanese Patent Publication Nos. 47-20635 and 50-22044, in which cationic polymers are used as a water-soluble polymer in order to enhance the adsorption of Mept compounds to the hair and head skin. However, the cationic polymers function as a coagulating agent for fine particulate Mept compounds having a specific size distribution, and produce considerable coagulation.

10 The method of forming colloid layer of cellulose-type water-soluble polymers for protection also has a drawback in that the cellulose polymers coagulate when they form a colloidal adsorption layer on a particle to be protected. For instance, addition of an aqueous solution of hydroxyalkylcellulose which is referred to in Japanese Patent Application Laid-open Nos. 53-14710, 53-14711, 53-97010 and 57-176906, results in losing the original size distribution, leading to losing the stable dispersability.

15 From this reason, conventional art was directed to make use of cellulose-type water-soluble polymers as a thickener to enhance the viscosity of shampoo base thereby allowing the incorporated fine particulate Mept compounds to settle more slowly, or further incorporating swelling clay or pearling agent in order to give structural viscosity to the system of the shampoo composition, thereby preventing settlement of Mept compounds. Such methods cannot avoid limitations to surfactants to be incorporated into a shampoo base or viscosity of the shampoo base.

20 Preparation of hair rinse compositions or hair conditioning compositions is also accompanied by the limitation to viscosity of the composition in the case where fine particulate Mept compounds are incorporated into the compositions. Namely, the bases are limited to a highly viscous ones such as of emulsion system or gel.

SUMMARY OF THE INVENTION

30 In view of the above, the present inventors have further made an earnest study in order to overcome such problems, and have found that when fine particulate Mept compounds are dispersed in water by aid of a specific dispersant, the resulting suspension is excellent in that it is highly stable under the ordinary storage conditions, is superior in resistivity to salts, and, even frozen, can be restored to the state before frozen if only heated to melt. Surprisingly, it was also found that when thus prepared aqueous dispersion of fine particulate Mept compounds was incorporated into a base of hair treatment compositions such as shampoos, hair rinses and hair lotions, the dispersion state of fine particulate Mept compounds could be maintained stably being free from limitations to storage conditions or to surfactants to be used in combination.

40 Accordingly, this invention provides an antimicrobial suspension comprising a dispersant and from 0.0015 to 60 wt.-% of a fine particulate polyvalent metal salt of 2-mercaptopyridine-N-oxide, characterized by having a size distribution in which particles having a size below 0.2 μm are contained in amounts not smaller than 50 wt.-%, particles having a size from 0.5 to 1.0 μm being 15 wt.-% or less and particles having a size over 1.0 μm being 2 wt.-% or less and the dispersant being selected from (A), (B) or (C):

45 (A) a polyglycol/polyamine condensation polymer, polyglycol/polyamine/alkylamine condensation polymer or alkyleneamine condensation polymer;

(B) at least one water-soluble polymer compound selected from hydroxyalkylcelluloses and partly quaternarized products thereof, and at least one non-ionic surfactant;

(C) at least one cationic polymer compound and at least one alkali metal salt, alkaline earth metal salt or aluminum salt of an inorganic acid.

50 Further, the present invention provides an antimicrobial hair treatment composition characterized by incorporating an antimicrobial suspension as defined in Claim 1.

BRIEF DESCRIPTION OF THE DRAWING

55 The sole figure is a graph showing relation between particle size of fine particulate Zpt and Amizet 5C.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

The Mept compounds according to the invention are represented by the following general formula:



10 wherein M represents a polyvalent metal atom and x represents the atomic value of M.

Examples of the polyvalent metal atom, M includes magnesium, barium, strontium, zinc, cadmium, tin and zirconium. Of these, zinc salt of 2-mercaptopyridine-N-oxide is preferred.

The fine particulate Mept compounds to be used in the present invention have a size distribution in which not smaller than 50 wt% of the particles have a size of 0.2 μ m (micron) or less. Preferably, the fine
15 particulate metal salts have such a size distribution that particles having a size ranging from 0.5 to 1.0 μ m (micron) are contained in amounts not larger than 15wt%, and particles having a size not smaller than 1.0 micron are contained 2wt% or less. The fine particulate Mept compounds having such size distribution can be prepared by any of the following methods:

(1) Method of utilizing a shear force of rigid body media such as glass beads having a particle size not
20 larger than 0.5 mm, with which Mept compounds are agitated thereby obtaining fine powder (Japanese patent Application No. 58-122845).

(2) Method of reacting a monovalent water-soluble salt of 2-mercaptopyridine-N-oxide and an water-soluble polyvalent metal salt in the presence of an water-soluble compound having a basic nitrogen atom in a molecular thereof under the condition of pH 3 to 7, or optionally, further proceeding this reaction in
25 the presence of a water-soluble compound having hydroxy group under a temperature of not higher than 0 $^{\circ}$ C (Japanese Patent Application No. 58-122846).

(3) Method of stirring a dispersion of Mept compounds together with rigid body media having a particle size not larger than 0.2 mm in the presence of a salt of (metha)acrylic acid-styrene sulfonic acid copolymer having an average molecular weight of 10,000 to 1,000,000 (Japanese patent Application No.
30 59-82690).

Among the dispersants usable in the present invention, the condensation polymers (A) are curable condensation products obtainable from firstly reacting a polyamine compound having from 2 to 10 carbon atoms with an ether of polyoxyalkylene glycol having a terminal halogen or hydroxyl group and having from 2 to 4 carbon atoms in an alkylene unit thereof, further reacting the resulting polyamine reaction product
35 having at least one hydrogen atom joined to a nitrogen atom with a bifunctional aliphatic compound having a functional group selected from epoxide and alpha-halo-beta-hydroxyalkyl. Alternatively, the thus obtained condensation product may further be reacted with amines having saturated or unsaturated hydrocarbon group having 10 to 24 carbon atoms thereby obtaining a curable condensation product.

Preferable condensation polymers are mentioned below.

40 (I) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000 and epichlorohydrin.

(II) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000, coconut oil fatty acid amine or beef tallow fatty acid amine and epichlorohydrin.

45 (III) Reaction product of dipropylenetriamine and bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600.

(IV) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600, coconut oil fatty acid amine or beef tallow fatty acid amine.

50 (V) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000 and bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600.

(VI) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600 and coconut oil fatty acid amine or beef tallow fatty acid amine.

55 (VII) Reaction product of diethylenetriamine, ethoxylated ethylene chlorohydrin and epichlorohydrin.

(VIII) Reaction product of diethylenetriamine, ethoxylated ethylene chlorohydrin, epichlorohydrin, and coconut oil fatty acid amine or beef tallow fatty acid amine.

(IX) Reaction product of dipropylenetriamine, ethoxylated glycerine chlorohydrin ether and epich-

lorohydrin.

(X) Reaction product of dipropylenetriamine, ethoxylated glycerine chlorohydrin ether, epichlorohydrin and coconut oil fatty acid amine or beef tallow fatty acid amine.

(XI) Reaction product of triethylenetetramine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000 and bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600.

(XII) Reaction product of triethylenetetramine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 1,000, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 600 and coconut oil fatty acid amine or beef tallow fatty acid amine.

(XIII) Reaction product of dipropylenetriamine and bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 200.

(XIV) Reaction product of dipropylenetriamine, bis-chlorohydrin ether of polyoxyethylene glycol having an average molecular weight of about 200 and coconut oil fatty acid amine or beef tallow fatty acid amine.


The ratio of the chlorine atoms to the hydrogen atoms of amino group in the reaction products of (I) to (XIV) is preferred to be in the range of 4:5 to 7:5.

Typical and preferable example of the condensation polymers is the product commercially sold under the trade name of Polyquart H from Henkel Co., Ltd., as a 50% aqueous solution.


Condensation polymers (A) are preferably incorporated in an amount of 1/100 to 10 times, more preferably 1/100 to 5 times of the fine particulate Mept compounds on the weight basis, and the balance is preferably an aqueous medium such as water or lower alcohol/water.

In the case where dispersant (A) is used, fine particulate Mept compounds should be incorporated in an amount of 0.0015 to 60%, and more preferably 0.1 to 50% of the total amount.

Examples of hydroxyalkyl celluloses of dispersant (B) of the invention include hydroxyethyl cellulose or hydroxypropyl cellulose. Cationic cellulose derivatives obtainable by quaternarizing the hydroxyalkyl celluloses are preferably those represented by the following formula:



wherein A_1 represents a residue of anhydroglucose unit, l represents an integer of from 50 to 20,000, and each R_1 represents a substitutional group represented by the following general formula (2):



wherein R' , R'' : alkylene group having 2 or 3 carbon atoms,

m : integer of from 0 to 10

n : integer of from 0 to 3

p : integer of from 0 to 10

R''' : alkylene group or hydroxyalkylene group having 1 to 3 carbon group,

R'_1 , R'_2 , R'_3 : same or different with each other and independently represent alkyl having up to 10 carbon atoms, aryl or aralkyl group, and may form a heterocyclic ring containing a nitrogen atom of the formula (2).


X : anion (chlorine, bromine, iodine, sulfuric acid, sulfonic acid, methylsulfuric acid, phosphoric acid, nitric acid.)

The cation substitution degree of the cationic celluloses preferably ranges from 0.01 to 1, in other words, average value of n per anhydroglucose unit is preferably 0.01 to 1, and more preferably, 0.02 to 0.5.

Average value of $m + p$ ranges from 1 to 3. Substitution degree of 0.01 or less is insufficient. On the other hand, that of 1 or more is objectionable in view of the yield of the reaction. Molecular weight of the cationic celluloses usable in the present invention ranges from about 100,000 to 300,000.

Specific examples of the non-ionic surfactants of dispersant (B) are mentioned below.


- (1) Polyoxyethylene alkyl or alkenyl ethers having an alkyl or alkenyl group having 10 to 20 carbon atoms on average and polyoxyethylene group having 1 to 20 units of ethylene oxide.
- (2) Polyoxyethylene alkylphenyl ethers having an alkyl group having 6 to 12 carbon atoms on average and polyoxyethylene group having 1 to 20 units of ethylene oxide.
- (3) Polyoxypropylene alkyl or alkenyl ethers having an alkyl or alkenyl group having 10 to 20 carbon atoms on average and polyoxypropylene group having 1 to 20 units of propylene oxide.
- (4) Polyoxybutylene alkyl or alkenyl ethers having an alkyl or alkenyl group having 10 to 20 carbon atoms on average and polyoxybutylene group having 1 to 20 units of butylene oxide.
- (5) Nonionic surface active agents having an alkyl or alkenyl group having 10 to 20 carbon atoms on average and added with 1 to 30 moles, in total, of ethylene oxide and propylene oxide or ethylene oxide and butylene oxide (a ratio of ethylene oxide and propylene oxide or butylene oxide is in the range of 0.1/9.9 to 9.9/0.1).
- (6) Higher fatty acid alkanolamides of the following formula or alkylene oxide adducts thereof




wherein R₂ represents an alkyl or alkenyl group having 7 to 21 carbon atoms, R₃ is a hydrogen atom or a methyl group, n₂ is an integer of 1 to 20, and m₃ is an integer of 0 to 20, and hardened castor oil.

Examples of the cationic polymer compounds of dispersant (C) in the invention include the following compounds (a), (b) and (c), and they are used solely or in combination of two or more.

(a) Copolymer-type cationic polymer compounds of dimethyldiallylammonium halide and acryl amide:




or



wherein R_{25} and R_{26} are the same or different with each other and independently represent a hydrogen atom or methyl, X represents a halogen atom, and l and m_4 are such integers that the sum of l and m_4 ranges from 180 to 2,000.

Although the compounds are represented by formula (a) here, they are not necessarily a block-type copolymer, and arrangement of the monomers may be arbitrary changed. Of these, compounds in which X is chlorine are sold under the tradename "Merquat 550" (Merck & Co., Inc./ U.S.A.)


(b) Polycondensation products of adipic acid and dialkylaminohydroxypropyl diethylenetriamine represented by the following formula and quaternarized products thereof:



wherein R_{27} and R_{28} are the same or different with each other and independently represent an alkyl group having from 1 to 18 carbon atoms and n_4 represents such integer that will give the molecular weight ranging from 100 to 100,000.

Of these, those whose R_{27} and R_{28} are methyl are available from Sandoz Co., Ltd. under the tradename of "Cartaretin F".

(c) Copolymers represented by the following formula and quaternarized products thereof:



wherein R_{29} represents a hydrogen atom, methyl group or ethyl group, Z represents an oxygen atom or

a group -NH-, R_{30} represents an alkylene group having 1 to 4 carbon atoms, R_{31} and R_{32} are the same or different with each other and independently represent an alkyl group having 1 to 18 carbon atoms, and p and q are such integers that will give a molecular weight ranging from 1,000 to 5,000,000.

Although the compounds are represented by the above formula, they are not necessarily a block-type copolymer, and arrangement of the monomers may be changed arbitrarily. Of these, compounds in which Z is oxygen, R_{29} is methyl, R_{30} is ethylene, R_{31} and R_{32} are methyl and quaternarized by $(C_2H_5)_2SO_4$ are available from GAF Corporation under the tradenames of "Gafqurt 755" and "Gafqurt 734".

The inorganic salts usable as a member of dispersant (C) include alkali metal salts, alkaline earth metal salts or aluminum salts of an inorganic acid such as chloric acid, sulfuric acid or nitric acid. Of these inorganic salts are preferred potassium sulfate, sodium sulfate, magnesium sulfate, aluminum sulfate, potassium nitrate, sodium nitrate, magnesium nitrate, calcium nitrate, aluminum nitrate, potassium chloride, sodium chloride, magnesium chloride, calcium chloride, aluminum chloride, potassium carbonate, sodium carbonate, aluminum carbonate. Among them, especially preferred ones are sodium sulfate, potassium nitrate, sodium nitrate, potassium chloride and sodium chloride.

The antimicrobial suspensions of the invention may be prepared according to conventional method by adding a powder of fine particulate Mept compounds or suspension thereof to a solution of the above-mentioned dispersants, but preferably be prepared by the following process.

Namely, the suspension is prepared by a method in which a powder of fine particulate Mept compounds is added to a 50% solution of dispersant (A) or a diluted solution thereof and agitated to give a uniform mixture, or by a method in which a dispersion of fine particulate Mept compounds is added to a 50% solution of dispersant (A) or a diluted solution thereof under agitation and further agitated to obtain a uniform mixture. The mixing is carried out using ordinary mixers such as propeller agitator apparatus, homogenizer mixers and sand mills.

When dispersant (B) is used, the following four ingredients are processed as described below.

- 1) 1 to 10% (as effective amount) of fine particulate Mept compounds,
- 2) Water-soluble polymer compound in an amount of 5 wt% or more of the fine particulate Mept compounds,
- 3) Non-ionic surfactant in an amount of 5 wt% or more of the water-soluble polymer compound,
- 4) ion-exchanged water as balance.

First, 2) is dissolved in 4), then added with 3), followed by further addition of 1) under agitation until uniform mixture is obtained. Alternatively, 2) is dissolved in 4), then added with 1) and mixed well, followed by further addition of 3) under agitation until uniform suspension is obtained. The mixing apparatus usable in the process are ordinary ones as described before.

Further, in case where dispersant (C) is used for preparing an antimicrobial suspension, an aqueous solution comprising 0.1 to 60 wt%, preferably 1 to 50 wt% of a cationic polymer compound and 0.1 to 25 wt%, preferably 1 to 25 wt% of an inorganic salt should firstly be prepared. Examples of solvents are water or mixture of water and lower alcohol. Into the thus obtained aqueous solution, a dispersion comprising 0.1 to 60 wt%, preferably 0.1 to 50 wt% of fine particulate Mept compounds is added and agitated until uniform mixture is obtained. The mixing is carried out by a similar apparatus described before. The thus obtained antimicrobial suspensions can be used as it is. Optionally, they may be subjected to filtration under pressure, by which step, Mept particles having improved surface are obtainable.

Examples of antimicrobial hair treatment compositions according to the second invention include compositions for shampoos, hair rinses or hair lotions. These compositions are prepared by incorporating the above described antimicrobial suspensions into a hair treatment base under agitation and uniformly mixing them. The antimicrobial suspension should be incorporated in such an amount that the amount of fine particulate Mept compounds are 0.01 to 10wt%, especially 0.05 to 5wt% of the total composition. The bases for hair treatments are those ordinarily used for these purposes. Among the hair treatment compositions, shampoo or hair rinse compositions are preferable in the practice of the invention. The bases for shampoos include anionic surfactants, amphoteric surfactants, cationic surfactants and mixtures thereof.

Specific examples of these surfactants are mentioned below.

Anionic surfactants:

- (1) Linear or branched alkylbenzenesulfonates having an alkyl group having 10 to 16 carbon atoms on average.
- (2) Polyoxyalkylene alkyl ether sulfates having a linear or branched alkyl group having 8 to 20 carbon atoms on average and added with 0.5 to 8 moles of ethylene oxide and/or propylene oxide in one molecule thereof on average.
- (3) Alkylsulfates having a linear or branched alkyl group having 10 to 20 carbon atoms on average.
- (4) Olefinsulfonates having 10 to 20 carbon atoms in one molecule thereof on average.


(5) Alkanesulfonates having 10 to 20 carbon atoms in one molecule thereof on average.

(6) Fatty acid salts having a linear or branched, saturated or unsaturated hydrocarbon chain having 10 to 20 carbon atoms on average.



(7) Alkylethoxycarboxylates having a linear or branched alkyl group having 10 to 20 carbon atoms and added with 0.5 to 8 moles of ethylene oxide in one molecule thereof on average.

(8) Alkyl or alkenylsuccinates having an alkyl or alkenyl group having 6 to 20 carbon atoms on average and partially neutralized salts thereof.

(9) Phosphate active agents of the formula




in which A represents




(in which R_1 represents a linear or branched, saturated or unsaturated hydrocarbon group, R_2 represents a hydrogen atom or a methyl group, m is an integer of from 0 to 6, and n is an integer of 1 to 6), B represents $-\text{OX}_2$ or A , and X_2 and X_2 independently represent a hydrogen atom or counter ion.

(10) Amino acid surface active agent of the formulae



in which R_3 represents an alkyl or alkenyl group having 7 to 21 carbon atoms, X_3 , X_4 and X_5 independently represent a hydrogen atom or counter ion.

(11) Acylated polypeptide surface active agents of the formula




in which R_4 represents an alkyl or alkenyl group having 7 to 21 carbon atoms, R_5 , R_6 and R_7 independently represent a side chain of an amino acid, n_1 is an integer of 1 to 6, and X_6 represents a hydrogen atom or counter ion.

The counter ions represented by X_1 to X_6 of these anionic surfactants generally include ions of alkali metals such as sodium or potassium; alkaline earth metals such as magnesium; ammonium ion and alkanolamine bases having 1 to 3 alkanol groups having 2 or 3 carbon atoms such as, for example,

monoethanolamine, diethanolamine, triethanolamine or triisopropanolamine.



Amphoteric surfactants:

(12) Alkylamine oxides (I) and amidoamine oxides (II) of the following formulae




in which R_8 represents an alkyl or alkenyl group having 10 to 20 carbon atoms, R_9 and R_{10} independently represent an alkyl group having 1 to 3 carbon atoms, and m_1 is an integer of 1 to 4.

(13) Alkyl or sulfobetaines (III) and amido or amidosulfobetaines (IV) of the following formulae




in which R_{11} and R_{12} independently represent an alkyl group having 1 to 4 carbon atoms, m_2 is an integer of 1 to 3, X_7 represents a $-\text{COO}^{\ominus}$ or SO_3^{\ominus} group, m_1 and R_8 have the same meanings as defined before, respectively.

(14) Imidazoline amphoteric surface active agents of the following formula




in which R_{13} is an aliphatic acid residue having 10 to 20 carbon atoms on the average, R_{14} represents sodium, hydrogen or $-\text{CH}_2\text{COOM}_2$, Z_1 represents $-\text{COOM}_2$, $-\text{CH}_2\text{COOM}_2$ or




in which M_2 represents a sodium-, hydrogen -atom or an organic base, X_8 represents a hydroxyl group, an acidic salt, an anionic surface active sulfate or a sulfated product.

(15) Amidoamine amphoteric surface active agents of the formula




in which R_{15} represents an alkyl or alkenyl group having 6 to 20 carbon atoms, R_{16} represents a hydrogen atom, $-\text{C}_2\text{H}_4\text{OH}$ or $-\text{C}_2\text{H}_4\text{OC}_2\text{H}_4\text{COOX}_9$, R_{17} represents $-\text{C}_2\text{H}_4\text{OH}$, $-\text{C}_2\text{H}_4\text{OC}_2\text{H}_4\text{COOX}_9$ or $-\text{C}_2\text{H}_4\text{COOX}_9$, and R_{18} represents a hydrogen atom or $-\text{C}_2\text{H}_4\text{COOX}_9$, X_9 represents a hydrogen-, alkali metal atom ammonium or organic ammonium. Cationic surfactants:

(16)




in which at least one of R_{20} , R_{21} , R_{22} and R_{23} represents an alkyl or alkenyl group having 8 to 24 carbon atoms and the other represent an alkyl group having 1 to 5 carbon atoms, and X' represents a halogen atom.

(17)



in which R_{20} , R_{21} , R_{22} and X' have the same meanings as defined before, respectively.

(18)



in which R_{20} , R_{21} and X' have the same meanings as defined before, respectively, n_3 is an integer of 1 to 20, and R_{24} represents an alkylene group having 2 to 3 carbon atoms.

It is to be noted that dispersant (B) requires a non-ionic surfactant as an essential component thereof, but when other dispersants are used, non-ionic surfactants may be incorporated as an arbitrary ingredient. Examples of the non-ionic surfactants include, aside from the before-mentioned polyox-

yethylene alkyl or alkenyl ethers, polyoxyethylene alkylphenyl ethers, polyoxypropylene alkyl or alkenyl ethers, polyoxybutylene alkyl or alkenyl ethers, non-ionic surfactants obtainable by adding propylene oxide or butylene oxide to ethylene oxide, and higher fatty acid alkanolamides or alkylene oxide adducts thereof, the following compounds are mentioned.

5 (19) Sucrose fatty acid esters obtainable from fatty acid having 10 to 20 carbon atoms on average and sucrose,

(20) Fatty acid glycerine monoester obtainable from fatty acid having 10 to 20 carbon atoms on average and glycerine.

Of these surfactants, agents (1), (2), (3), (4), (5), (6), (11), (I) and (II) of (12), (13), (14), (15),
10 polyoxyethylene alkyl or alkenyl ethers and higher fatty acid alkanolamides or alkyleneoxide adducts thereof are preferred. These surfactants may be used singly or in combination. Suitable amount of the surfactants is, for example, in the range not less than 5wt%, preferably from 10 to 40wt% in total, of the composition.

The antimicrobial hair treatment compositions may further comprise, aside from the above-described essential ingredients, any arbitrary ingredients ordinarily used for these purposes. Examples of such
15 arbitrary ingredients include: solubilizers such as propylene glycol, glycerine or urea; viscosity modifiers such as ethyl alcohol, isopropyl alcohol, methyl cellulose or higher alcohols; preservatives; antiseborrheic agents; keratin-soluble or swelling substances such as sulfur, salicylic acid and enzymes; deodorants; pearling agents; lotionizing agents; perfumes; colorants; UV absorbers; antioxidants; preservatives.

20 Action:

The function of the dispersants to be used in the present invention is not clear but it is considered that the dispersants make the particle surface of the fine particulate Mept compounds improved, thereby aggregation of the particles is prevented.

25

Effects of the Invention:

Several shampoos and hair rinses containing the Mept compounds are put on the market. However, in order to stably incorporate the Mept compounds into shampoo or hair rinse compositions, it is unavoidable
30 to make them highly viscous or to add specific types of polymers or clay minerals. This leads to the disadvantage that the compositions become poor in performance, e.g. with shampoos, the foaming performance becomes poor with objectionable texture of the hair after shampooing.

Different from conventional method of modifying the viscosity of dispersion medium for improving the dispersion stability of Mept compounds, the inventive method utilizes fine particulate Mept compounds,
35 thereby obtaining an antimicrobial suspension of very stable dispersion. Antimicrobial hair treatment compositions which incorporate fine particulate Mept compounds are excellent in storage stability and in feel on use.

Examples:

40

The present invention is described by way of referential examples and examples.

The size distribution of Zpt in the referential examples or examples was determined by a centrifugal automatic size distribution measuring instrument CAPA-500 (available from K.K. Horiba Seisakusho). Water was used as a medium for dispersion. Density of the composition, viscosity and the specific gravity of Zpt
45 particles were 1, 0.8mPas (cps) (30 °C), and 1.78, respectively.

Referential Example 1

0.29 g of hepta hydride salt of zinc sulfate and 99.3 g of mixture of 1% solution of Polyquat H and
50 water/ethanol (65/35) were charged in a reaction vessel and cooled down to -25 °C, to which 0.75 g of aqueous 40% solution of sodium salt of 2-mercaptopyridine-N-oxide was charged at a time and reacted. After further 1 minute of agitation at the same temperature, the settled Zpt fine powder was collected. The size distribution of the fine particulate Zpt is shown in Table 1.

55

Table 1

	Particle Size (μ)	Distribution (wt%)	Cumulative Total (wt%)
5	1.00 <	0.0	0.0
10	1.00 - 0.95	0.0	0.0
	0.95 - 0.90	0.6	0.6
	0.90 - 0.85	0.0	0.6
	0.85 - 0.80	0.0	0.6
15	0.80 - 0.75	0.2	0.8
	0.75 - 0.70	1.1	1.9
	0.70 - 0.65	0.0	1.9
20	0.65 - 0.60	0.0	1.9
	0.60 - 0.55	8.4	2.3
	0.55 - 0.50	0.3	2.5
	0.50 - 0.45	0.0	2.5
25	0.45 - 0.40	0.2	2.7
	0.40 - 0.35	1.4	4.1
	0.35 - 0.30	2.1	6.2
	0.30 - 0.25	1.4	7.7
30	0.25 - 0.20	1.6	9.2
	0.20 - 0.15	2.7	11.9
	0.15 - 0.10	8.7	20.6
35	0.10 - 0.05	76.1	96.7
	0.05 - 0.00	3.3	100.0
	Average Size	0.08 μ	

40

Referential Example 2

45 70 ml of a commercially available Zpt dispersion (commercial product A having active ingredient of 50 wt%) and 130 ml of glass beads (media) having a size of 44 to 63 microns were mixed (media/dispersion ratio by volume = 65/35) and placed in a 400 ml sand grinder (made by Igarashi Machine Manufacturing Co., Ltd.). Grinder discs were rotated for 3 hours at a peripheral speed of 5 m/sec. The temperature within the grinder was 20 to 25°C. The resulting dispersion was filtered under pressure to obtain 31 ml of fine
50 particulate Zpt. The size distribution of thus obtained fine particulate Zpt is shown in Table 2.

55

Table 2

	Particle Size (μ)	Distribution (wt%)	Cumulative Total (wt%)
5	0.90 <	0.0	0.0
10	0.90 - 0.84	0.0	0.0
	0.84 - 0.78	0.0	0.0
15	0.78 - 0.72	1.4	1.4
	0.72 - 0.66	3.3	4.7
	0.66 - 0.60	0.0	4.7
20	0.60 - 0.54	2.4	7.1
	0.54 - 0.48	1.9	9.0
	0.48 - 0.42	2.8	11.8
25	0.42 - 0.36	4.3	16.1
	0.36 - 0.30	5.7	21.8
	0.30 - 0.24	7.3	29.1
30	0.24 - 0.18	12.5	41.6
	0.18 - 0.12	22.2	63.8
	0.12 - 0.06	32.9	96.7
35	0.06 - 0.00	3.3	100
40	Average Size	0.16 μ	


45

Referential Example 3

23 g of commercially available Zpt powder, 4.5 g of a water-soluble copolymer salt of the formula having a molecular weight of 680,000,

50

55



63 g of water, and 187 g of glass beads (media) having a size of 0.1 to 0.2 mm were mixed (media/dispersion ratio by volume = 63/37) and placed in a 400 ml sand grinder (made by Igarashi Machine Manufacturing Co., Ltd.). In the grinder, discs were rotated for 12 hours at a peripheral speed of 6 m/sec. The temperature in the sand grinder was found to be 20 to 25 °C. The content was filtered under pressure to obtain 40 g of fine particulate Zpt dispersion. When the media are washed twice each with 70 g of water, 98% of Zpt could be recovered. The size distribution of thus obtained fine particulate Zpt is shown in Table 3.

Table 3

	Particle Size (μ)	Distribution (wt%)	Cumulative Total (wt%)
5	0.90 <	0.0	0.0
10	0.90 - 0.84	0.0	0.0
	0.84 - 0.78	0.0	0.0
15	0.78 - 0.72	0.0	0.0
	0.72 - 0.66	0.0	0.0
	0.66 - 0.60	0.0	0.0
20	0.60 - 0.54	0.8	0.8
	0.54 - 0.48	0.0	0.8
	0.48 - 0.42	1.0	1.8
25	0.42 - 0.36	0.1	1.9
	0.36 - 0.30	1.4	3.3
	0.30 - 0.24	4.2	7.5
30	0.24 - 0.18	3.0	10.5
	0.18 - 0.12	7.4	17.9
	0.12 - 0.06	20.2	38.2
35	0.06 - 0.00	61.8	100
40	Average Size	0.05 μ	

45

Example 1

Antimicrobial suspensions as shown in Table 1 were prepared by adding a certain amount of suspension containing Zpt in an amount of 50% to an aqueous solution of a water-soluble polymer of a predetermined concentration, and agitating the mixture. The appearance of the suspensions were observed immediately after the preparation, after 10 days storage at 25°C, after addition of sodium chloride (final concentration = 3%, after 10 days of storage), and after frozen and re-melted (from -20°C to room temperature) by naked eyes. The results are shown in Table 4. In the table, symbols denote the following meanings.

- 55 o: uniform dispersion
 F: flocculation
 S: sedimentation
 A: irreversible aggregation

Table 4
(wt %)

Antimicrobial Suspension	Inventive Products										Comparative Products									
	1	2	3	4	5	20	40	60	80	10	20	30	40	50	60	70	80	90	10	
Fine Particulate 2pt Suspension (active ingredient 50%) ¹⁾																				
Commercially Sold 2pt Suspension A (active ingredient 50%) ²⁾																				
Commercially Sold 2pt Suspension B (active ingredient 48%) ³⁾																				
Polyglycol/Polyamine Polycondensation Product (active ingredient 50%) ⁴⁾																				
Cationic Polymer Solution C (active ingredient 30%) ⁵⁾																				
" D (active ingredient 40%) ⁶⁾																				
" E (active ingredient 19%) ⁷⁾																				
Hydroxyethyl Cellulose ⁸⁾																				
Carboxyvinyl Polymel ⁹⁾																				
Water																				
Appearance: Immediately after the preparation																				
After storage for 10 days at 25° C																				
Ten days after addition of 3% sodium chloride																				
After melting of frozen suspension																				

1) Product of Referential Example 2

2) Size Distribution

	Particle Size (μ)	Distribution (wt%)	Cumulative Total (wt%)
5	2.00 <	0.0	0.0
	2.00 - 1.90	0.0	0.0
	1.90 - 1.80	0.6	0.6
	1.80 - 1.70	7.7	8.3
10	1.70 - 1.60	4.1	12.4
	1.60 - 1.50	0.6	13.0
	1.50 - 1.40	0.0	13.0
15	1.40 - 1.30	0.0	13.0
	1.30 - 1.20	0.6	13.6
20	1.20 - 1.10	2.8	16.4
	1.10 - 1.00	0.0	16.4
	1.00 - 0.90	1.9	18.3
25	0.90 - 0.80	9.0	27.3
	0.80 - 0.70	1.4	28.7
	0.70 - 0.60	4.6	33.3
30	0.60 - 0.50	9.8	43.1
	0.50 - 0.40	11.9	55.0
	0.40 - 0.30	16.1	71.1
35	0.30 - 0.20	17.8	88.9
	0.20 - 0.10	9.5	98.4
40	0.10 - 0.00	1.6	100.0
	Average Size	0.44 μ	

45 3) Size Distribution

50

55

	Particle Size (μ)	Distribution (wt%)	Cumulative Total (wt%)
5	3.00 <	0.0	0.0
	3.00 - 2.85	0.0	0.0
	2.85 - 2.70	2.1	2.1
10	2.70 - 2.55	1.3	3.4
	2.55 - 2.40	0.0	3.4
	2.40 - 2.25	1.9	5.3
	2.25 - 2.10	3.1	8.4
15	2.10 - 1.95	2.7	11.1
	1.95 - 1.80	1.5	12.6
	1.80 - 1.65	1.0	13.6
	1.65 - 1.50	1.5	15.1
20	1.50 - 1.35	9.6	24.7
	1.35 - 1.20	14.4	39.1
	1.20 - 1.05	5.6	44.7
25	1.05 - 0.90	6.3	51.0
	0.90 - 0.75	7.9	58.9
	0.75 - 0.60	9.0	67.9
	0.60 - 0.45	11.9	79.8
30	0.45 - 0.30	12.1	91.9
	0.30 - 0.15	7.5	99.4
	0.15 - 0.00	0.6	100.0
35	Average Size	0.91 μ	

40 4) Polyquart H (Henkel Co., Ltd.)

5) Polycondensation product of Adipic acid/Dimethylamino hydroxypropane diethylenetriamine (Cartaretin F, Sandoz Co., Ltd.)

6) 1:1 Copolymer of methyldivinylimidazolium/vinylpyrrolidone (Luviquat FC 550, BASF Co., Ltd.)

7) Copolymer of vinylpyrrolidone/dimethylaminoethylmethacrylate (Gafquat 755N, GAF Co., Ltd.)

45 8) Average polymerization 1550

9) Carbopol 941 (Goodrich Co., Ltd.)

As shown in Table 4, only inventive products incorporating fine particulate Zpt and condensation polymerization product of polyglycol/polyamine were excellent in lasting uniform suspension without any flocculation or sedimentation taken place under any condition.

50

Example 2

A surfactant was added to water and dissolved to prepare a uniform solution having a final concentration referred to in Table 5. An antimicrobial suspension prepared in Example 1 was added to the resulting solution under agitation, further added with perfume and colorant. The pH was adjusted to 7 by citric acid, and the viscosity to 800 cps by ethanol, thereby obtaining shampoo compositions shown in Table 2. The appearance of the compositions were observed after 30 days storage at room temperature, at 50 °C and at 40 °C, respectively, and after frozen and re-melted (from -20 °C to room temperature) by naked eyes. The

results are shown in Table 5. The symbols have the same meanings as indicated in Example 1.

As shown in Table 5, only inventive shampoo compositions incorporating fine particulate Zpt and condensation polymerization product of polyglycol/polyamine were excellent in lasting stable suspension under any condition.

Table 5

Shampoo Composition		(wt %)															
		Inventive Products					Comparative Products										
		5	6	7	8	9	10	11	12	13	14	15					
Suspension of Antimicrobial Agent (Inventive Product 2, Zpt cont. 10%) (Comparative Product 1, Zpt cont. 5%)		10	10	10	10	20	5										
(Comparative Product 3, Zpt cont. 10%)							10										
"							10										
"							10										
"							10										
"							10										
"							10										
"							10										
Sodium Polyoxyethylene(2.5)lauryl ether Sulfate (25%)		60	40	40	40	40	47										
Triethanolamine Lauryl Sulfate (40%)							30										
Sodium C ₁₄ ,16- α -Olefin sulfonate (30%)							15										
Imidazoline-type Amphoteric Surfactant (30%)							20										
Coconut Oil Fatty Acid Amidobetaine							4										
Coconut Oil Fatty Acid Acyl-L-glutamic Triethanolamine							6										
Sodium Lauroyl Sarcosinate							3										
Coconut Oil Fatty Acid Diethanolamide							3										
Water and other additives							27										
Appearance: Immediately after the preparation		○	○	○	○	○	○	○	○	○	○	○	○	○	○		
After 30 days storage at Room Temperature		○	○	○	○	○	○	○	○	○	○	○	○	○	○		
After 30 days storage at 50 °C		○	○	○	○	○	○	○	○	○	○	○	○	○	○		
After 30 days storage at 40 °C		○	○	○	○	○	○	○	○	○	○	○	○	○	○		
After melting of frozen suspension		○	○	○	○	○	○	○	○	○	○	○	○	○	○		

Example 3

5	Hair Rinse Composition:		
	1) Distearyltrimethylammonium chloride	2%	
	2) Cetyl alcohol	2	
10	3) Propylene glycol	3	
	4) Antimicrobial suspension (Inventive product 2, Zpt 10%)	3	
15	5) Perfume	0.5	
	6) Colorant	small amount	
	7) Citric acid	small amount	
20	8) Water	balance	

25 In a uniform solution of (1) and (8) was uniformly dispersed (4) and heated, followed by adding a hot uniform solution of (2) and (3) under agitation, cooling and adding (5), (6) and (7) to obtain a hair rinse composition. Thus obtained hair rinse composition was found to keep good and stable suspension over a long period without being flocculated or settled after heated or frozen and re-melted.

Example 4

30

Anti-dandruff Lotion:

35	Fine particulate Zpt (50%) 4% (obtained in Referential Example 2)		
40	Polyquart H	2.0	
	Propylene glycol	5	
	Ethanol	10	
45	Perfume	small amount	
	Water	balance	

50 The procedure of Example 1 was followed to prepare an anti-dandruff lotion. The lotion was found to be stable over 1 month at room temperature.

On the contrary, when no Polyquart H was used, Zpt was recognized to settle in 3 days.

Example 5

55

Aqueous Antimicrobial:

5	Fine particulate Zpt (50%) (obtained in Referential Example 2)	0.2%
	Polyquart H	0.2
10	Water	suitable amount
	<hr/>	
	Total	100

15 An aqueous antimicrobial was obtained in the same manner as in Example 2. The antimicrobial was stable over 1 month and showed good antimicrobial effects when applied to trees.

Example 6

20 An antimicrobial dispersion of fine particulate Zpt of the following formulation was prepared, then diluted by ion-exchanged water and determined the size distribution by the centrifugal precipitation method. The results are shown in the sole Figure.

(Formulation)

	Hydroxyethylcellulose *1	1.2%
30	Zpt dispersion of Referential Example 2 (active ingredient 50 wt%)	8
	Polyoxyethylene coconut oil fatty acid monoethanolamide *2	0.1 to 0.8
35	Ion-exchanged water	balance
	(Note) *1: HEC UNICEL QP 4400H, product of Daicel Chemical Industry Co., Ltd.	
40	*2: Amizet 5C, product of Kawaken Fine Chemical Co., Ltd.	

45

(Preparation)

50 Hydroxyethylcellulose was added to ion-exchanged water and heated to dissolve, followed by cooling down to room temperature, adding polyoxyethylene coconut fatty acid monoethanolamide to obtain a uniform mixture. Dispersion of fine particulate Zpt according to Referential Example 2 was added to the resulting mixture, and further agitated to obtain a uniform mixture.

Example 7

55

Shampoo Composition:**(Formulation)**

EP 0 173 259 B1

5	1) Antimicrobial suspension according to Example 6 (containing Amizet 5C by 0.6%)	25%
	2) Laurylsulfate triethanolamine salt (active ingredient 40%)	40
10	3) Lauric acid diethanolamide	3
	4) Phosphoric acid	suitable amount
	5) Colorant	small amount
15	6) Perfume	0.5%
	7) Ethanol	suitable amount
20	8) Ion-exchanged water	balance

(Preparation)

25 In a uniform solution of 2) and 8) was dispersed 1) under agitation, and 3), 5) and 6) were further added under agitation until uniform mixture was obtained. 4) and 7) were added to the resulting solution in a suitable amount to adjust pH to 7, and viscosity to 200 cp (B-viscometer), thereby obtaining an anti-dandruff shampoo composition. This composition showed good dispersion after storage for 2 weeks at 40° C.

30 Example 8

Hair Rinse Composition:

(Formulation)

35	1) Antimicrobial suspension according to Example 6 (containing Amizet 5C by 0.6%)	12.5%
40	2) Stearyltrimethylammonium chloride	3
	3) Polyoxyethylene cetyl ether (EO 5)	2
45	4) Liquid paraffin	0.5
	5) Isostearic acid	0.5
50	6) Perfume	0.5
	7) Colorant	small amount
	8) Ion-exchanged water	balance

65

(Preparation)

In a uniform solution of 2) and 8) was dispersed 1) under agitation until uniformly mixed, and 3), 4), 5), 6) and 7) were further added and mixed uniformly to prepare an anti-dandruff rinse composition. This composition showed good dispersion after storage for 2 weeks at 40° C.

5 **Examples 9 - 16**

Shampoo compositions shown in Table 6 were prepared following the methods of Examples 6 and 7. These products showed good dispersion after storage for 2 weeks at 40° C.

10 **Comparative Example**

Comparative products 16 to 20 shown in Table 6 were prepared. Products which were not treated by water-soluble polymer/non-ionic surfactant were flocculated and sedimented. Commercially sold Zpt treated by water-soluble polymer/non-ionic surfactant showed sedimentation. The fine particulate Zpt which was not
15 treated by non-ionic surfactant but was treated only by water-soluble polymer showed sedimentation. The fine particulate Zpt treated by methylcellulose/non-ionic surfactant showed sedimentation. The fine particulate Zpt treated by carboxymethylcellulose/non-ionic surfactant was flocculated and sedimented.

20

25

30

35

40

45

50

55

Table 6

	Inventive Products										Comparative Products					
	11	12	13	14	15	16	17	18	18	17	18	18	19	20	19	20
Antimicrobial Suspension *3																
Zpt Dispersion of Referential Example 2 (active ingredient 50%)	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Commercially sold Zpt Dispersion (active ingredient 50%, 0.45 μm)	0.2	0.2		0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2				
Hydroxyethyl Cellulose *4																
Hydroxypropyl Cellulose *5		0.2														
Cationic Cellulose *6			0.2													
Methyl Cellulose *7																
Carboxymethyl Cellulose *8																
Polyoxyethylene(5) Coconut Oil Fatty Acid Monothanolamide *9	0.1	0.1	0.1		0.1	0.1	0.1	0.1	0.1		0.1	0.1	0.1	0.1	0.1	0.1
Polyoxyethylene(80) Hardened Castor Oil *10				0.1												
Secondary Alcohol E.O. Additive (ED90) *11					0.1											
Sodium Polyoxyethylene(2.5)laurylether Sulfate (active ingredient 25%)	60	60	60	60	60		40	40	60	60	60	60	60	60	60	60
Sodium C14,18-α-olefinsulfonate (active ingredient 30%)	3	3	3	3	3	50					3	3	3	3	3	3
Coconut Oil Fatty Acid Diethanolamide																
Sodium Lauroyl Sarcosinate																
Coconut Oil Fatty Acid Monosodium Acyl- L-glutamate																
Perfume							0.5									
Colorant							small amount									
pH modifier							adjust pH to 7									
Ethanol							suitable amount to adjust viscosity of the shampoo composition to 200cp.									
Water							balance									
Stability after storage for 2 weeks at 40°C ±12	0	0	0	0	0	0	0	0	0	0	0	F	S	S	S	F

- *3: Prepared according to Example 6.
- *4: HEC UNICEL QP4400H (product of Daicel Chemical Industry
Co., Ltd.)
- *5: HPC-M (product of Sanyo-Kokusaku Pulp Co., Ltd.)
- *6: Polymer JR 400
- *7: Metholose 60 SH (product of Shin-Etsu Chemical Co., Ltd.)
- *8: Daicel CMC #1260 (product of Daicel Chemical Industry Co.,
Ltd.)
- *9: Amizet 5C (product of Kawaken Fine Chemical Co., Ltd.)
- *10: HCO-80 (product of Nikko Chemical Co., Ltd.)
- *11: Softanol 90 (product of Nippon Shokubai K.K.)
- *12: Dispersion State

Uniform dispersion: O

Sedimentation: S

Flocculation: F

Example 17

58 g of water, 15 g of sodium chloride and 17 g of Cartaretin F (active ingredient 30%) were mixed up, followed by adding 10 g of fine particulate Zpt (active ingredient 50%) under agitation, and subjected to further agitation to obtain a suspension of fine particulate Zpt.

20 parts by weight (hereinafter simply referred to as parts) of the resulting suspension, which was in gel state, were added to 80 parts of sodium alkylether sulfate (active ingredient 2.5%) under agitation, and further mixed well, thereby obtaining a uniform suspension. It was found to contain particles having as same size as measured prior to the treatment.

Zpt suspensions shown in Table 7 were prepared in the same manner as described above. Appearance of the suspension was observed after mixed with sodium alkylether sulfate under the same condition as mentioned above, and after 40 days storage at 25°C, respectively. In the Table, "O", "A", "F" and "S" denote the following meanings respectively.

- O: Uniform suspension
A: Irreversible aggregation
F: Flocculation
S: Sedimentation

As shown in Table 7, inventive products 19 to 25 which make use of the compound selected from polycondensation product of adipic acid-dimethylaminohydroxypropyl diethylenetriamine, copolymerization product of dimethylaminoethyl methacrylate-vinylpyrrolidone and copolymerization product of dimethyldiallylammoniumchloride-acrylamide, inorganic salt and fine particulate Zpt gave excellent and uniform dispersion while keeping the size of fine particulate Zpt even in the presence of surfactants, and were not sedimented or irreversibly aggregated. On the contrary, comparative products 21 to 27 generated flocculation or sedimentation, and were not agreeable.

Table 7

21: Cartaretin F (Sandoz Co.)	22: Gafquat 755 (GAF Corp.)	23: Gafquat 734 (GAF Corp.)	24: Merquat (Merck & Co., Inc.)
-------------------------------	-----------------------------	-----------------------------	---------------------------------

55

27

Table 8
(wt %)

Zpt Suspension No.		Inventive Products													
Component	26	27	28	28	30	31	32	33	34	35	36	37	38	39	40
Suspension of fine particulate Zpt according to Referential Example 2 (active ingredient 50%, average size 0.16μ)	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Merquat 550	3.75	3.75	5	5	3.75	5	3.75	3.75	5	5	3.75	5	5	6.25	6.25
Gafquat 734															
Gafquat 755															
Cartaretin F															
Sodium Chloride	7.5	7.5	10	10	10	10	7.5	7.5	10	10	10	7.5	10	7.5	10
Water	Balance														
Appearance: 80 parts of Sodium Polyoxyethylene(2.5) Lauryl Sulfate is added to 20 parts of the above suspension and stirred.	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
Appearance: After 40 days storage at 25° C	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○

Example 19

Shampoo compositions shown in Table 9 were prepared making use of Zpt suspensions obtained in Example 17.

(preparation)

A surfactant, aqueous solution of a surfactant, and water were uniformly mixed, to which Zpt suspension prepared according to Example 17 was added under agitation, followed by further agitation until uniform mixture was obtained. The resulting mixture was added with perfume and colorant, adjusted pH to 7 by citric acid, and viscosity to 1,000 - 500 by ethanol.

*1: necessary amount for adjusting pH to 7

*2: necessary amount for modifying the viscosity in a range from 1,000 to 500

*3: Miranol C2M (Miranol Co. Ltd.)

10 *4: Rebon 2000 (Sanyo Kasei K.K.)

As shown in Table 9, inventive shampoo products 41 to 48 gave excellent suspension even after 2 month storages at room temperature, 40 °C, and 50 °C without being sedimented or flocculated. Further, after frozen and re-melted, the inventive products were found to restore good suspension. On the contrary, comparative product 28 sedimented without flocculation, and comparative products 29 and 30 sedimented with flocculation.

20

25

30

35

40

45

50

55

Table 9

(wt %)

Component	Shampoo Composition No.										Inventive Products								Comparative Products			
	41	42	43	44	45	46	47	48	28	29	30	41	42	43	44	45	46	47	48	28	29	30
Suspension of Inventive Product 20	20																					
Suspension of Inventive Product 24		20																				
Suspension of Inventive Product 25			20																			
Suspension of Comparative Product 24																						
Suspension of Comparative Product 26																						
Suspension of fine particulate Zpt (active ingredient 50%)																						
Sodium Polyoxyethylene(2.5) Lauryl Sulfate (active ingredient 25%)																						
Sodium C ₁₄ -18- α -Olefin sulfonate (active ingredient 30%)																						
Coconut Oil Fatty Acid Diethanolamide																						
Imidazoline-type Amphoteric Surfactant (active ingredient 40%) *3																						
Coconut Oil Fatty Acid Amidobetaine (active ingredient 30%) *4																						
Sodium Lauroyl Sarcosinate																						
Monosodium Coconut Oil Fatty Acid Acyl-L-glutamate																						
Perfume																						
Colorant																						
Citric Acid																						
Ethanol																						
Water																						
Appearance: After storage for 2 months at Room Temperature																						
"																						
"																						
"																						
After melting of frozen suspension																						

Example 20

Hair rinse Composition:

(Formulation)

	Stearyltrimethyl ammonium chloride	2 wt%
	Cetyl alcohol	3
5	Propylene glycol	3
	Zpt suspension (Inventive Product 20)	5
	Perfume	0.5
10	Colorant	small amount
	Citric acid	small amount
15	Water	balance

Into a uniform solution of stearyltrimethyl ammoniumchloride and water was added and suspended a suspension of Inventive product 20 and heated, to which a uniform solution of cetyl alcohol and propylene glycol was added under further agitation. After cooling, perfume, colorant and citric acid were added to prepare a hair rinse composition.


Thus obtained hair rinse compositions gave good suspension stability at room temperature, at 40° C, and at 50° C.

25 Claims


1. An antimicrobial suspension comprising a dispersant and from 0.0015 to 60 wt.-% of a fine particulate polyvalent metal salt of 2-mercaptopyridine-N-oxide, characterized by having a size distribution in which particles having a size below 0.2 μm are contained in amounts not smaller than 50 wt.-%, particles having a size from 0.5 to 1.0 μm being 15 wt.-% or less and particles having a size over 1.0 μm being 2 wt.-% or less and the dispersant being selected from (A), (B) or (C):
 - (A) a polyglycol/polyamine condensation polymer, polyglycol/polyamine/alkylamine condensation polymer or alkyleneamine condensation polymer;
 - (B) at least one water-soluble polymer compound selected from hydroxyalkylcelluloses and partly quaternarized products thereof, and at least one non-ionic surfactant;
 - (C) at least one cationic polymer compound and at least one alkali metal salt, alkaline earth metal salt or aluminum salt of an inorganic acid.
2. The antimicrobial suspension according to Claim 1, wherein said polyvalent metal salt of 2-mercaptopyridine-N-oxide is a zinc salt.
3. The antimicrobial suspension according to Claim 1, wherein said polyvalent metal salt of 2-mercaptopyridine-N-oxide having a size distribution in which particles having a size below 0.2 μm are contained in amounts not smaller than 50 wt% is incorporated in said suspension by 0.0015 to 60 wt%, and said dispersant (A) is incorporated in said suspension by 1/100 to 10 times of the amount of said fine particulate polyvalent metal salt of 2-mercaptopyridine-N-oxide.
4. The antimicrobial suspension according to Claim 1, wherein said at least one water-soluble polymer compound of dispersant (B) is selected from a hydroxyethylcellulose, a hydroxypropylcellulose, and a reaction product of a hydroxyethylcellulose or a hydroxypropylcellulose with glycidyltrimethylammonium chloride.
5. The antimicrobial suspension according to Claim 1, wherein said at least one non-ionic surfactant of dispersant (B) is selected from polyoxyalkylenealkyl or polyoxyalkylene alkenyl ethers having straight or branched alkyl or alkylene group having 10 to 20 carbon atoms on average and polyoxyalkylene group having 1 to 20 units of alkylene oxide having 2 to 4 carbon atoms, polyoxyalkylene alkylphenyl ethers, alkyleneoxide adducts of higher aliphatic acid alkanolamide and hardened castor oil.

6. The antimicrobial suspension according to Claim 1, wherein said at least one cationic polymer compound of dispersant (C) is selected from the following compounds of (a) to (c):

(a) A copolymer-type cationic polymer compound of dimethyl diallylammonium halide and acrylamide represented by either formula of




or




wherein, R_{25} and R_{26} are the same or different with each other and independently represent a hydrogen atom or methyl group, X represents a halogen atom, and t and m_4 are such integers that the sum of t and m_4 ranges from 180 to 2,000.

(b) A polycondensation product of adipic acid and dialkylaminohydroxypropyl diethylenetriamine represented by the following formula or quaternarized products thereof:



wherein, R_{27} and R_{28} are the same or different with each other, and independently represent an alkyl group having 1 to 18 carbon atoms, and n_4 is such an integer that gives a molecular weight of ranging from 100 to 100,000.

(c) A copolymer represented by the following formula or quaternarized products thereof;



20 wherein R_{29} represents a hydrogen atom, methyl group or ethyl group, Z represents an oxygen atom or a group -NH-, R_{30} represents an alkylene group having 1 to 4 carbon atoms, R_{31} and R_{32} are the same or different with each other and independently represent an alkyl group having 1 to 18 carbon atoms, and p and q independently represent such an integer that gives a molecular weight of ranging from 1,000 to 5,000,000.



- 25 7. An antimicrobial hair treatment composition characterized by incorporating an antimicrobial suspension as defined in claim 1.
8. The hair treatment composition according to Claim 7, wherein said composition is a shampoo composition or a hair rinse composition.
- 30 9. An antimicrobial hair treatment composition according to Claim 7 or Claim 8, wherein said hair treatment composition comprises a surfactant for shampoo use containing a mixture of one or more agents selected from anionic surfactants, amphoteric surfactants, non-ionic surfactants and cationic surfactants.
- 35 10. The antimicrobial hair treatment composition according to Claim 9, wherein said surfactant for shampoo use is contained in an amount of 5 wt% or more based on the total composition.

Revendications

- 40 1. Suspension antimicrobienne comprenant un dispersant et de 0,0015 à 60% en poids d'un sel particulaire, fin, de métal polyvalent du mercapto-2 pyridine-N-oxyde, caractérisé par une répartition granulométrique dans laquelle des particules ayant une taille inférieure à 0,2 μm sont présentes dans des quantités non inférieures à 50% en poids, les particules ayant une taille de 0,5 à 1,0 μm représentant 15% en poids ou moins, et les particules ayant une taille dépassant 1,0 μm représentant 2% en poids ou moins, et le dispersant étant choisi parmi (A), (B) ou (C) :
- 45 (A) un polymère de condensation polyglycol/polyamine, un polymère de condensation polyglycol/polyamine/alkylamine ou un polymère de condensation d'une alkylèneamine ;
- (B) au moins un composé polymère soluble dans l'eau choisi parmi les hydroxyalkylcelluloses et leurs produits partiellement quaternisés, et au moins un agent tensio-actif non-ionique ;
- 50 (C) au moins un composé polymère cationique et au moins un sel de métal alcalin, sel de métal alcalino-terreux ou sel d'aluminium d'un acide minéral.
2. Suspension antimicrobienne selon la revendication 1, dans laquelle ledit sel de métal polyvalent du mercapto-2 pyridine-N-oxyde est un sel de zinc.
- 55 3. Suspension antimicrobienne selon la revendication 1, dans laquelle ledit sel de métal polyvalent du mercapto-2 pyridine-N-oxyde ayant une répartition granulométrique dans laquelle des particules ayant

une dimension inférieure à 0,2 μm sont présentes dans des quantités non inférieures à 50% en poids, est incorporé dans ladite suspension à raison de 0,0015 à 60% en poids, et ledit dispersant (A) est incorporé dans ladite suspension à raison de 1/100 à 10 fois la quantité dudit sel particulière, fin, de métal polyvalent du mercapto-2 pyridine-N-oxyde.


- 5
4. Suspension antimicrobienne selon la revendication 1, dans laquelle ledit (ou lesdits) composé(s) polymère(s) soluble(s) dans l'eau du dispersant (B) est (sont) choisi(s) parmi une hydroxyéthylcellulose, une hydroxypropylcellulose, et un produit de réaction d'une hydroxyéthylcellulose ou d'une hydroxypropylcellulose avec le chlorure de glycidyl triméthyl ammonium.
- 10
5. Suspension antimicrobienne selon la revendication 1, dans laquelle ledit (ou lesdits) agent(s) tensio-actif(s) non-ionique(s) du dispersant (B) est (ou sont) choisi(s) parmi les polyoxyalkylène alkyl ou polyoxyalkylène alcényl éthers ayant un groupe alkyle ou alkylène à chaîne droite ou ramifié ayant 10 à 20 atomes de carbone en moyenne et un groupe polyoxyalkylène ayant 1 à 20 unités d'oxyde d'alkylène ayant 2 à 4 atomes de carbone, les polyoxyalkylène alkylphényl éthers, les produits d'addition d'oxyde d'alkylène d'alcanolamide d'acide aliphatique supérieur et d'huile de ricin durcie.
- 15
6. Suspension antimicrobienne selon la revendication 1, dans laquelle ledit (ou lesdits) composé(s) polymère(s) cationique(s) du dispersant (C) est (ou sont) choisi(s) parmi les composés suivants (a) à (c) :
- 20
- (a) un composé polymère cationique de type copolymère d'halogénure de diméthyl diallylammonium et d'acrylamide représenté par l'une ou l'autre des formules :



dans lesquelles :

- R_{25} et R_{26} sont identiques ou différents l'un de l'autre et représentent indépendamment un atome d'hydrogène ou un groupe méthyle ;
- X représente un atome d'halogène ; et
- l et m_1 sont des nombres entiers tels que la somme de l et m_1 se situe dans la plage de 180 à 2 000.


(b) un produit de polycondensation d'acide adipique et de dialkylaminohydroxypropyl diéthylène triamine représenté par la formule suivante, ou les produits quaternisés de celui-ci :



dans laquelle :

- R_{27} et R_{28} sont identiques ou différents l'un de l'autre, et représentent indépendamment un groupe alkyle ayant 1 à 18 atomes de carbone ; et
- n_4 est un nombre entier tel qu'il donne une masse moléculaire se situant dans la plage de 100 à 100 000.

(c) un copolymère représenté par la formule suivante, ou les produits quaternisés de celui-ci :



40 dans laquelle :

- R_{29} représente un atome d'hydrogène, un groupe méthyle ou un groupe éthyle ;
- Z représente un atome d'oxygène ou un groupe
- R_{30} représente un groupe alkylène ayant 1 à 4 atomes de carbone ;
- R_{31} et R_{32} sont identiques ou différents l'un de l'autre et représentent indépendamment un
- p et q représentent indépendamment un nombre entier tel qu'il donne une masse moléculaire se situant dans la plage de 1 000 à 5 000 000.

50 7. Composition de traitement capillaire antimicrobien caractérisée par l'incorporation d'une suspension antimicrobienne telle que définie à la revendication 1.

8. Composition de traitement capillaire selon la revendication 7, dans laquelle ladite composition est une composition de shampooing ou une composition de rinçage capillaire.


55 9. Composition de traitement capillaire antimicrobien selon la revendication 7 ou la revendication 8, dans laquelle ladite composition de traitement capillaire comprend un agent tensio-actif pour utilisation dans un shampooing, contenant un mélange d'un ou plusieurs agents choisis parmi les agents tensio-actifs anioniques, les agents tensio-actifs amphotères, les agents tensio-actifs non-ioniques et les agents

tensio-actifs cationiques.


10. Composition de traitement capillaire antimicrobien selon la revendication 9, dans laquelle ledit agent tensio-actif pour utilisation dans un shampooin, est présent dans une quantité de 5% en poids ou
5 davantage sur la base de la composition totale.

Patentansprüche

1. Antimikrobielle Suspension, umfassend ein Dispersionsmittel und von 0,0015 bis 60 Gew.% eines feinteiligen polyvalenten Metallsalzes von 2-Mercaptopyridin-N-oxid, gekennzeichnet durch eine Größenverteilung, bei der Teilchen mit einer Größe unter 0,2 μm in Mengen von nicht weniger als 50 Gew.% enthalten sind, Teilchen mit einer Größe von 0,5 - 1,0 μm 15 Gew.% oder weniger ausmachen und Teilchen mit einer Größe über 1,0 μm 2 Gew.% oder weniger ausmachen und wobei das Dispersionsmittel ausgewählt ist aus (A), (B) oder (C):
15 (A) ein Polyglycol/Polyamin-Kondensationspolymerisat, Polyglycol/Polyamin/Alkylamin-Kondensationspolymerisat oder Alkylamin Kondensationspolymerisat;
(B) mindestens eine wasserlöslichen Polymerverbindung, ausgewählt aus Hydroxyalkylcellulosen und teilweise quaternierten Produkten derselben, und mindestens ein nicht-ionisches Surfactans;
(C) mindestens eine kationische Polymerverbindung und mindestens ein Alkalimetallsalz, Erdalkalimetallsalz oder Aluminiumsalz einer anorganischen Säure.
2. Antimikrobielle Suspension gemäß Anspruch 1, wobei das polyvalente Metallsalz von 2-Mercaptopyridin-N-Oxid ein Zinksalz ist.
- 25 3. Antimikrobielle Suspension gemäß Anspruch 1, wobei das polyvalente Metallsalz von 2-Mercaptopyridin-N-Oxid, das eine Größenverteilung aufweist, bei der Teilchen mit einer Größe unter 0,2 μm in Mengen von nicht weniger als 50 Gew.% enthalten sind, der Suspension mit 0,0015 bis 60 Gew.% einverleibt ist und wobei das Dispersionsmittel (A) der Suspension mit 1/100 bis zum 10-fachen der Menge des feinteiligen polyvalenten Metallsalzes von 2-Mercaptopyridin-N-Oxid einverleibt ist.
- 30 4. Antimikrobielle Suspension, wobei die mindestens eine wasserlösliche Polymerverbindung von Dispersionsmittel (B) ausgewählt ist aus einer Hydroxyethylcellulose, einer Hydroxypropylcellulose und einem Reaktionsprodukt einer Hydroxyethylcellulose oder einer Hydroxypropylcellulose mit Glycidyltrimethylammoniumchlorid.
- 35 5. Antimikrobielle Suspension gemäß Anspruch 1, wobei das mindestens eine nicht-ionische Surfactans von Dispersionsmittel (B) ausgewählt ist aus Polyoxyalkylenalkyl- oder Polyoxyalkylenalkenyläthern mit graden oder verzweigten Alkyl- oder Alkylengruppen mit 10 bis 20 Kohlenstoffatomen im Durchschnitt und einer Polyoxyalkylengruppe mit 1 bis 20 Einheiten von Alkylenoxid mit 2 bis 4 Kohlenstoffatomen, Polyoxyalkylenalkylphenyläthern, Alkylenoxidaddukten von höherem aliphatischem Säurealkanolamid und gehärtetem Rizinusöl.
- 40 6. Antimikrobielle Suspension gemäß Anspruch 1, wobei die mindestens eine kationische Polymerverbindung von Dispersionsmittel (C) ausgewählt ist aus den folgenden Verbindungen (a) bis (c):
45 (a) eine kationische Polymerverbindung vom Copolymerisattyp von Dimethyldiallylammoniumhalogenid und von Acrylamid, dargestellt durch die folgenden Formeln




oder




15 wobei R_{25} und R_{26} gleich oder verschieden voneinander sind und unabhängig für ein Wasserstoffatom oder eine Methylgruppe stehen, X für ein Halogenatom steht und l und m_1 derartige ganze Zahlen sind, daß die Summe von l und m_1 im Bereich von 180 bis 2 000 liegt;

20 (b) ein Polykondensationsprodukt von Adipinsäure und Dialkylaminohydroxypropyl-diethylentriamin gemäß der folgenden Formel oder quaternisierte Produkte desselben:



35 wobei R_{27} und R_{28} gleich oder voneinander verschieden sind und unabhängig eine Alkylgruppe mit 1 bis 18 Kohlenstoffatomen bedeuten und n_1 eine derartige ganze Zahl ist, daß ein Molekulargewicht im Bereich von 100 bis 100 000 erhalten wird;

40 (c) ein Copolymerisat der folgenden Formel oder quaternisierte Produkte desselben:



wobei R_{29} ein Wasserstoffatom, eine Methylgruppe oder Ethylgruppe bedeutet, Z für ein Sauerstoff-

atom oder eine Gruppe -NH-, steht, R_{3c} eine Alkylengruppe mit 1 bis 4 Kohlenstoffatomen bedeutet, R_{31} und R_{32} gleich oder voneinander verschieden sind und unabhängig für eine Alkylgruppe mit 1 bis 18 Kohlenstoffatomen stehen und p und q unabhängig eine derartige ganze Zahl bedeuten, daß ein Molekulargewicht im Bereich von 1000 bis 5 000 000 erhalten wird.

5

7. Antimikrobielles Haarbehandlungsmittel, gekennzeichnet durch die Einverleibung einer Antimikrobiellen Suspension, wie sie in Anspruch 1 definiert ist.

10

8. Haarbehandlungsmittel gemäß Anspruch 7, wobei das Mittel als Schampoozusammensetzung oder als eine Haarspülzusammensetzung vorliegt.

15

9. Antimikrobielles Haarbehandlungsmittel gemäß Anspruch 7 oder 8, wobei das Haarbehandlungsmittel ein Surfactans für Schampooverwendungszwecke umfaßt, enthaltend eine Mischung von einem oder mehreren Mitteln, ausgewählt aus anionischen Surfactantien, amphoterischen Surfactantien, nicht-ionischen Surfactantien und kationischen Surfactantien.

20

10. Antimikrobielles Haarbehandlungsmittel gemäß Anspruch 9, wobei das Surfactans für Schampooverwendungszwecke in einer Menge von 5 Gew.% oder mehr, bezogen auf die Gesamtzusammensetzung, enthalten ist.

25

30

35

40

45

50

55

FIGURE

